

Studying the exotic decay of $^{70}\text{Kr} \rightarrow ^{70}\text{Br}$

Monday, 17 August 2020 17:25 (15 minutes)

The β -decay of ^{70}Kr can be used to test the predictions of different theoretical models. First of all, the effects of $T = 0$ pn-pairing can be investigated through the study of the decay rates [1]. Furthermore the theoretical strength distributions of the β -decay of ^{70}Kr show clear differences depending on the shape of the ground state, hence assumptions on the shape of the ground state can be tested [2]. Finally ^{70}Kr might lay on the rp-process path and accordingly its half-life might play a role in rp-process network calculations [3]. Despite its importance, the knowledge on the β -decay of this isotope is rather limited. The half-life is known only with 15% accuracy, and only one γ -transition was identified to date [4,5].

To make the tests of these models possible, we've conducted an experiment at RIKEN-RIBF. The nucleus of interest was produced using a ^{78}Kr primary beam with a kinetic energy of 345 MeV/nucleon and intensity of about 300 pnA. The fragments were separated by BigRIPS using in-flight method, then were stopped in the WAS3ABi active silicon detector [6]. The β -delayed γ -rays were detected by the EURICA cluster array surrounding the implantation station [7].

The experiment significantly increased our knowledge on the structure of the daughter nucleus populated in the β -decay. A precise half-life value, with an uncertainty in the order of 2%, have been derived from time correlations of the β -particles and the decay-curves of the observed γ -transitions. Furthermore, after the identification of several new γ -transitions, a detailed level scheme – including about 20 transitions – have been derived [8]. The experimental approach, as well as the new half-life value and the level scheme, will be presented, along with an insight on the theoretical interpretations of the experimental results.

- [1] A. L. Goodman, Phys. Rev. C 60, 014311 (1999)
- [2] P. Sarriguren, Phys. Rev. C 83, 025801 (2011)
- [3] H. Schatz, et al., Phys. Rev. Lett. 86, 3471 (2001)
- [4] M. Oinonen, et al., Phys. Rev. C 61, 035801 (2000)
- [5] G. de Angelis et al., Eur. Phys. J. A 12, 51 (2001)
- [6] S. Nishimura, Prog. Theor. Exp. Phys. 03C006 (2012)
- [7] P.-A. Söderström, et al., Nucl. Instrum. Meth. B 317, 649 (2013)
- [8] A. Vitéz-Sveiczler et al., Acta Phys. Pol. B 51, 587 (2020)

Field of your work

Experiential nuclear physics

Primary authors: VITEZ-SVEICZER, Andras (Institute for Nuclear Research (Atomki)); Dr ALGORA, Alejandro (IFIC, CSIC-University); Dr ANABEL ISABEL, Morales (IFIC, CSIC-University); Prof. RUBIO, Berta (IFIC, CSIC-University); Dr KISS, Gabor Gyula (Institute for Nuclear Research (Atomki))

Presenter: VITEZ-SVEICZER, Andras (Institute for Nuclear Research (Atomki))

Session Classification: Young Scientist Session 1