

## Direct measurement of the cross section for $^{102}\text{Pd}(p,g)^{103}\text{Ag}$ reaction in the p-process

Thursday, 31 August 2023 14:00 (5 minutes)

The study of the p-process is of paramount importance in unraveling the origin of heavy elements in the universe. To describe the entire p-nuclei nucleosynthesis process, a comprehensive reaction network involving over ten thousand nuclear reactions is required, and accurate measurements of some key reaction cross sections are essential for determining reaction rates.  $^{102}\text{Pd}$  is one of the more than 30 p-nuclei, and the  $^{102}\text{Pd}(p,g)^{103}\text{Ag}$  reaction is one of its significant destruction reactions. Experimental studies for the p-nucleus  $^{102}\text{Pd}$  indicate that the reaction rate for  $^{102}\text{Pd}(p,g)^{103}\text{Ag}$  is significantly higher than HF predictions. There are significant discrepancies in the available data on the  $^{102}\text{Pd}(p,g)^{103}\text{Ag}$  reaction cross section in the low-energy regime relevant to nuclear astrophysics. In light of these discrepancies, a direct measurement was carried out to determine the reaction cross section of  $^{102}\text{Pd}(p,g)^{103}\text{Ag}$  within the energy range of 1.9-2.8 MeV. The measurement was conducted utilizing the 2\*1.7 MV tandem accelerator at China Institute of Atomic Energy (CIAE). The latest cross section data were obtained using offline activation measurement technique based on the low background anti-muon and anti-Compton spectrometer in CIAE.

The latest results have extended the cross section of  $^{102}\text{Pd}(p,g)^{103}\text{Ag}$  to the lowest energy range of proton down to 1.9 MeV. The newly measured cross section data provide valuable experimental references for the calculation of statistical models, particularly in the low-energy regime of interest in nuclear astrophysics. These results contribute to a better understanding of the p-process and its implications for the nucleosynthesis of heavy elements in the universe.

**Primary authors:** LIU, Fulong (Center for Nuclear Study, the University of Tokyo); Prof. GUO, Bing (China Institute of Atomic Energy); Prof. HE, Chuangye (China Institute of Atomic Energy); Dr CHENG, Hao (China Institute of Atomic Energy); Dr BO, Nan (China Institute of Atomic Energy)

**Presenter:** LIU, Fulong (Center for Nuclear Study, the University of Tokyo)

**Session Classification:** Poster Session